

A3.2: Number of patients with grade 0-4 intestinal LENT SOMA by toxicity at 2 weeks

| | | | Control N=28 | | HBO N=50 | | Total N=78 | |
|--------------|-------------------------------|---------|-----------------|-------|-------------|--------|---------------|-------|
| | | | N | % | N | % | N | % |
| Subjective | Stool frequency | 0 | 6 | 21.4% | 13 | 26.0% | 19 | 24.4% |
| | | 1 | 18 | 64.3% | 26 | 52.0% | 44 | 56.4% |
| | | 2 | 4 | 14.3% | 9 | 18.0% | 13 | 16.7% |
| | | 3 | 0 | 0.0% | 2 | 4.0% | 2 | 2.6% |
| | | | | | | | | |
| | Stool consistency | 0 | 25 | 89.3% | 42 | 84.0% | 67 | 85.9% |
| | | 1 | 0 | 0.0% | 2 | 4.0% | 2 | 2.6% |
| | | 2 | 3 | 10.7% | 5 | 10.0% | 8 | 10.3% |
| | | 3 | 0 | 0.0% | 1 | 2.0% | 1 | 1.3% |
| | | | | | | | | |
| | Pain | 0 | 16 | 57.1% | 22 | 44.0% | 38 | 48.7% |
| | | 1 | 4 | 14.3% | 10 | 20.0% | 14 | 17.9% |
| | | 2 | 5 | 17.9% | 14 | 28.0% | 19 | 24.4% |
| | | 3 | 3 | 10.7% | 3 | 6.0% | 6 | 7.7% |
| | | 4 | 0 | 0.0% | 1 | 2.0% | 1 | 1.3% |
| Constipation | 0 | 27 | 96.4% | 49 | 98.0% | 76 | 97.4% | |
| | 1 | 1 | 3.6% | 1 | 2.0% | 2 | 2.6% | |
| Objective | Melena | 0 | 26 | 92.9% | 47 | 94.0% | 73 | 93.6% |
| | | 1 | 2 | 7.1% | 3 | 6.0% | 5 | 6.4% |
| | Weight loss | 0 | 26 | 92.9% | 49 | 98.0% | 75 | 96.2% |
| | | Missing | 2 | 7.1% | 1 | 2.0% | 3 | 3.8% |
| | Stricture | 0 | 25 | 89.3% | 49 | 98.0% | 74 | 94.9% |
| | | Missing | 3 | 10.7% | 1 | 2.0% | 4 | 5.1% |
| | Ulceration | 0 | 25 | 89.3% | 49 | 98.0% | 74 | 94.9% |
| | | Missing | 3 | 10.7% | 1 | 2.0% | 4 | 5.1% |
| Management | Pain | 0 | 22 | 78.6% | 35 | 70.0% | 57 | 73.1% |
| | | 1 | 3 | 10.7% | 11 | 22.0% | 14 | 17.9% |
| | | 2 | 3 | 10.7% | 3 | 6.0% | 6 | 7.7% |
| | | 3 | 0 | 0.0% | 1 | 2.0% | 1 | 1.3% |
| | | | | | | | | |
| | Stool consistency / frequency | 0 | 10 | 35.7% | 12 | 24.0% | 22 | 28.2% |
| | | 1 | 11 | 39.3% | 13 | 26.0% | 24 | 30.8% |
| | | 2 | 6 | 21.4% | 24 | 48.0% | 30 | 38.5% |
| | | 3 | 1 | 3.6% | 1 | 2.0% | 2 | 2.6% |
| | Bleeding | 0 | 25 | 89.3% | 48 | 96.0% | 73 | 93.6% |
| | | 1 | 3 | 10.7% | 2 | 4.0% | 5 | 6.4% |
| | Stricture | 0 | 25 | 89.3% | 50 | 100.0% | 75 | 96.2% |
| | | 1 | 1 | 3.6% | 0 | 0.0% | 1 | 1.3% |
| | | Missing | 2 | 7.1% | 0 | 0.0% | 2 | 2.6% |
| | Ulceration | 0 | 26 | 92.9% | 50 | 100.0% | 76 | 97.4% |
| Missing | | 2 | 7.1% | 0 | 0.0% | 2 | 2.6% | |

A3.3: Number of patients with grade 0-4 intestinal LENT SOMA by toxicity at 12 months

| | | | Control N=26 | | HBO N=46 | | Total N=72 | | |
|-------------|-------------------------------|---------|-----------------|--------|-------------|--------|---------------|--------|-------|
| | | | N | % | N | % | N | % | |
| Subjective | Stool frequency | 0 | 7 | 26.9% | 8 | 17.4% | 15 | 20.8% | |
| | | 1 | 12 | 46.2% | 24 | 52.2% | 36 | 50.0% | |
| | | 2 | 5 | 19.2% | 11 | 23.9% | 16 | 22.2% | |
| | | 3 | 2 | 7.7% | 3 | 6.5% | 5 | 6.9% | |
| | Stool consistency | 0 | 22 | 84.6% | 32 | 69.6% | 54 | 75.0% | |
| | | 1 | 1 | 3.8% | 3 | 6.5% | 4 | 5.6% | |
| | | 2 | 3 | 11.5% | 11 | 23.9% | 14 | 19.4% | |
| | Pain | 0 | 14 | 53.8% | 20 | 43.5% | 34 | 47.2% | |
| | | 1 | 3 | 11.5% | 6 | 13.0% | 9 | 12.5% | |
| | | 2 | 6 | 23.1% | 15 | 32.6% | 21 | 29.2% | |
| | | 3 | 2 | 7.7% | 5 | 10.9% | 7 | 9.7% | |
| | | 4 | 1 | 3.8% | 0 | 0.0% | 1 | 1.4% | |
| | Constipation | 0 | 26 | 100.0% | 46 | 100.0% | 72 | 100.0% | |
| | Objective | Melena | 0 | 25 | 96.2% | 45 | 97.8% | 70 | 97.2% |
| | | | 1 | 0 | 0.0% | 1 | 2.2% | 1 | 1.4% |
| 2 | | | 1 | 3.8% | 0 | 0.0% | 1 | 1.4% | |
| Weight loss | | 0 | 26 | 100.0% | 46 | 100.0% | 72 | 100.0% | |
| Stricture | | 0 | 26 | 100.0% | 45 | 97.8% | 71 | 98.6% | |
| | | Missing | 0 | 0.0% | 1 | 2.2% | 1 | 1.4% | |
| Ulceration | | 0 | 26 | 100.0% | 45 | 97.8% | 71 | 98.6% | |
| | Missing | 0 | 0.0% | 1 | 2.2% | 1 | 1.4% | | |
| Management | Pain | 0 | 22 | 84.6% | 36 | 78.3% | 58 | 80.6% | |
| | | 1 | 2 | 7.7% | 6 | 13.0% | 8 | 11.1% | |
| | | 2 | 1 | 3.8% | 2 | 4.3% | 3 | 4.2% | |
| | | 3 | 1 | 3.8% | 2 | 4.3% | 3 | 4.2% | |
| | Stool consistency / frequency | 0 | 11 | 42.3% | 19 | 41.3% | 30 | 41.7% | |
| | | 1 | 10 | 38.5% | 8 | 17.4% | 18 | 25.0% | |
| | | 2 | 5 | 19.2% | 18 | 39.1% | 23 | 31.9% | |
| | | 3 | 0 | 0.0% | 1 | 2.2% | 1 | 1.4% | |
| | Bleeding | 0 | 26 | 100.0% | 44 | 95.7% | 70 | 97.2% | |
| | | 1 | 0 | 0.0% | 2 | 4.3% | 2 | 2.8% | |
| | Stricture | 0 | 26 | 100.0% | 46 | 100.0% | 72 | 100.0% | |
| | Ulceration | 0 | 26 | 100.0% | 46 | 100.0% | 72 | 100.0% | |

A3.4 Maximum grade reported for each scale at baseline

| Maximum grade at baseline | | Control N=29 | | HBO N=55 | | Total N=84 | |
|---------------------------|---|-----------------|-------|-------------|-------|---------------|-------|
| | | N | % | N | % | N | % |
| Subjective | 0 | 2 | 6.9% | 3 | 5.5% | 5 | 6.0% |
| | 1 | 13 | 44.8% | 10 | 18.2% | 23 | 27.4% |
| | 2 | 10 | 34.5% | 28 | 50.9% | 38 | 45.2% |
| | 3 | 4 | 13.8% | 14 | 25.5% | 18 | 21.4% |
| Objective | 0 | 23 | 79.3% | 45 | 81.8% | 68 | 81.0% |
| | 1 | 3 | 10.3% | 4 | 7.3% | 7 | 8.3% |
| | 2 | 2 | 6.9% | 5 | 9.1% | 7 | 8.3% |
| | 3 | 1 | 3.4% | 1 | 1.8% | 2 | 2.4% |
| Management | 0 | 8 | 27.6% | 12 | 21.8% | 20 | 23.8% |
| | 1 | 8 | 27.6% | 16 | 29.1% | 24 | 28.6% |
| | 2 | 11 | 37.9% | 23 | 41.8% | 34 | 40.5% |
| | 3 | 2 | 6.9% | 4 | 7.3% | 6 | 7.1% |

A3.5 Maximum grade reported for each scale at 2 weeks

| Maximum grade at 2 weeks | | Control N=28 | | HBO N=50 | | Total N=78 | |
|--------------------------|---------|-----------------|-------|-------------|-------|---------------|-------|
| | | N | % | N | % | N | % |
| Subjective | 0 | 5 | 17.9% | 7 | 14.0% | 12 | 15.4% |
| | 1 | 11 | 39.3% | 17 | 34.0% | 28 | 35.9% |
| | 2 | 9 | 32.1% | 20 | 40.0% | 29 | 37.2% |
| | 3 | 3 | 10.7% | 5 | 10.0% | 8 | 10.3% |
| | 4 | 0 | 0.0% | 1 | 2.0% | 1 | 1.3% |
| Objective | 0 | 23 | 82.1% | 46 | 92.0% | 69 | 88.5% |
| | 1 | 2 | 7.1% | 3 | 6.0% | 5 | 6.4% |
| | Missing | 3 | 10.7% | 1 | 2.0% | 4 | 5.1% |
| Management | 0 | 6 | 21.4% | 11 | 22.0% | 17 | 21.8% |
| | 1 | 12 | 42.9% | 14 | 28.0% | 26 | 33.3% |
| | 2 | 7 | 25.0% | 23 | 46.0% | 30 | 38.5% |
| | 3 | 1 | 3.6% | 2 | 4.0% | 3 | 3.8% |
| | Missing | 2 | 7.1% | 0 | 0.0% | 2 | 2.6% |

A3.6 Maximum grade reported for each scale at 12 months

| Maximum grade at 12 months | | Control N=26 | | HBO N=46 | | Total N=72 | |
|----------------------------|---------|-----------------|-------|-------------|-------|---------------|-------|
| | | N | % | N | % | N | % |
| Subjective | 0 | 6 | 23.1% | 5 | 10.9% | 11 | 15.3% |
| | 1 | 9 | 34.6% | 16 | 34.8% | 25 | 34.7% |
| | 2 | 7 | 26.9% | 19 | 41.3% | 26 | 36.1% |
| | 3 | 3 | 11.5% | 6 | 13.0% | 9 | 12.5% |
| | 4 | 1 | 3.8% | 0 | 0.0% | 1 | 1.4% |
| Objective | 0 | 25 | 96.2% | 44 | 95.7% | 69 | 95.8% |
| | 1 | 0 | 0.0% | 1 | 2.2% | 1 | 1.4% |
| | 2 | 1 | 3.8% | 0 | 0.0% | 1 | 1.4% |
| | Missing | 0 | 0.0% | 1 | 2.2% | 1 | 1.4% |
| Management | 0 | 10 | 38.5% | 17 | 37.0% | 27 | 37.5% |
| | 1 | 9 | 34.6% | 9 | 19.6% | 18 | 25.0% |
| | 2 | 6 | 23.1% | 18 | 39.1% | 24 | 33.3% |
| | 3 | 1 | 3.8% | 2 | 4.3% | 3 | 4.2% |

A3.7 Maximum overall intestinal LENT SOMA grade reported at baseline

| Maximum score at baseline | | Control N=29 | | HBO N=55 | | Total N=84 | |
|---------------------------|---|-----------------|-------|-------------|-------|---------------|-------|
| | | N | % | N | % | N | % |
| | 0 | 1 | 3.4% | 3 | 5.5% | 4 | 4.8% |
| | 1 | 9 | 31.0% | 5 | 9.1% | 14 | 16.7% |
| | 2 | 13 | 44.8% | 31 | 56.4% | 44 | 52.4% |
| | 3 | 6 | 20.7% | 16 | 29.1% | 22 | 26.2% |

A3.8 Maximum overall intestinal LENT SOMA grade reported at 2 weeks

| Maximum score at 2 weeks | | Control N=28 | | HBO N=50 | | Total N=78 | |
|--------------------------|---|-----------------|-------|-------------|-------|---------------|-------|
| | | N | % | N | % | N | % |
| | 0 | 3 | 10.7% | 6 | 12.0% | 9 | 11.5% |
| | 1 | 12 | 42.9% | 14 | 28.0% | 26 | 33.3% |
| | 2 | 9 | 32.1% | 23 | 46.0% | 32 | 41.0% |
| | 3 | 4 | 14.3% | 6 | 12.0% | 10 | 12.8% |
| | 4 | 0 | 0.0% | 1 | 2.0% | 1 | 1.3% |

A3.9 Maximum overall intestinal LENT SOMA grade reported at 12 months

| Maximum score at 12 months | Control N=26 | | HBO N=46 | | Total N=72 | |
|-------------------------------|-----------------|-------|-------------|-------|---------------|-------|
| | N | % | N | % | N | % |
| 0 | 5 | 19.2% | 4 | 8.7% | 9 | 12.5% |
| 1 | 9 | 34.6% | 13 | 28.3% | 22 | 30.6% |
| 2 | 8 | 30.8% | 23 | 50.0% | 31 | 43.1% |
| 3 | 3 | 11.5% | 6 | 13.0% | 9 | 12.5% |
| 4 | 1 | 3.8% | 0 | 0.0% | 1 | 1.4% |

NOT FOR CITATION

Appendix 4 CTCAE

The number and proportion of patients with each grade of each toxicity are shown split by treatment group at each time point

A4.1 Baseline CTCAE toxicities

| | | Control N=29 | | HBO N=54 | | Total N=83 | |
|---------------------|---|-----------------|-------|-------------|-------|---------------|-------|
| | | N | % | N | % | N | % |
| Abdominal pain | 0 | 15 | 53.6% | 21 | 38.9% | 36 | 43.4% |
| | 1 | 10 | 35.7% | 16 | 29.6% | 26 | 31.3% |
| | 2 | 1 | 3.6% | 10 | 18.5% | 11 | 13.3% |
| | 3 | 3 | 10.7% | 7 | 13.0% | 10 | 12.0% |
| Bloating | 0 | 17 | 60.7% | 26 | 48.1% | 43 | 51.8% |
| | 1 | 8 | 28.6% | 17 | 31.5% | 25 | 30.1% |
| | 2 | 4 | 14.3% | 11 | 20.4% | 15 | 18.1% |
| Constipation | 0 | 26 | 92.9% | 49 | 90.7% | 75 | 90.4% |
| | 1 | 3 | 10.7% | 4 | 7.4% | 7 | 8.4% |
| | 2 | 0 | 0.0% | 1 | 1.9% | 1 | 1.2% |
| Diarrhoea | 0 | 21 | 75.0% | 39 | 72.2% | 60 | 72.3% |
| | 1 | 1 | 3.6% | 1 | 1.9% | 2 | 2.4% |
| | 2 | 6 | 21.4% | 12 | 22.2% | 18 | 21.7% |
| | 3 | 1 | 3.6% | 1 | 1.9% | 2 | 2.4% |
| | 4 | 0 | 0.0% | 1 | 1.9% | 1 | 1.2% |
| Faecal incontinence | 0 | 9 | 32.1% | 18 | 33.3% | 27 | 32.5% |
| | 1 | 6 | 21.4% | 19 | 35.2% | 25 | 30.1% |
| | 2 | 14 | 50.0% | 16 | 29.6% | 30 | 36.1% |
| | 3 | 0 | 0.0% | 1 | 1.9% | 1 | 1.2% |
| Flatulence | 0 | 9 | 32.1% | 20 | 37.0% | 29 | 34.9% |
| | 1 | 11 | 39.3% | 25 | 46.3% | 36 | 43.4% |
| | 2 | 9 | 32.1% | 9 | 16.7% | 18 | 21.7% |
| Frequency | 0 | 11 | 39.3% | 11 | 20.4% | 22 | 26.5% |
| | 1 | 6 | 21.4% | 17 | 31.5% | 23 | 27.7% |
| | 2 | 11 | 39.3% | 26 | 48.1% | 37 | 44.6% |
| | 3 | 1 | 3.6% | 0 | 0.0% | 1 | 1.2% |
| Rectal bleeding | 0 | 10 | 35.7% | 27 | 50.0% | 37 | 44.6% |
| | 1 | 7 | 25.0% | 6 | 11.1% | 13 | 15.7% |
| | 2 | 11 | 39.3% | 16 | 29.6% | 27 | 32.5% |
| | 3 | 1 | 3.6% | 5 | 9.3% | 6 | 7.2% |
| Rectal pain | 0 | 22 | 78.6% | 48 | 88.9% | 70 | 84.3% |
| | 1 | 4 | 14.3% | 2 | 3.7% | 6 | 7.2% |
| | 2 | 2 | 7.1% | 4 | 7.4% | 6 | 7.2% |
| | 3 | 1 | 3.6% | 0 | 0.0% | 1 | 1.2% |
| Tenesmus | 0 | 6 | 21.4% | 17 | 31.5% | 23 | 27.7% |
| | 1 | 13 | 46.4% | 16 | 29.6% | 29 | 34.9% |
| | 2 | 10 | 35.7% | 21 | 38.9% | 31 | 37.3% |
| Urgency | 0 | 3 | 10.7% | 5 | 9.3% | 8 | 9.6% |
| | 1 | 3 | 10.7% | 10 | 18.5% | 13 | 15.7% |
| | 2 | 21 | 75.0% | 37 | 68.5% | 58 | 69.9% |
| | 3 | 2 | 7.1% | 2 | 3.7% | 4 | 4.8% |
| GI disorders | 0 | 27 | 96.4% | 47 | 87.0% | 74 | 89.2% |
| | 1 | 1 | 3.6% | 6 | 11.1% | 7 | 8.4% |
| | 2 | 1 | 3.6% | 1 | 1.9% | 2 | 2.4% |

A4.2 2 week CTCAE toxicities

| | | HBO N=50 | | Control N=28 | | Total N=78 | |
|---------------------|---|-------------|-------|-----------------|-------|---------------|-------|
| | | N | % | N | % | N | % |
| Abdominal pain | 0 | 27 | 54.0% | 15 | 53.6% | 42 | 53.8% |
| | 1 | 12 | 24.0% | 8 | 28.6% | 20 | 25.6% |
| | 2 | 7 | 14.0% | 3 | 10.7% | 10 | 12.8% |
| | 3 | 4 | 8.0% | 2 | 7.1% | 6 | 7.7% |
| Bloating | 0 | 27 | 54.0% | 17 | 60.7% | 44 | 56.4% |
| | 1 | 20 | 40.0% | 6 | 21.4% | 26 | 33.3% |
| | 2 | 3 | 6.0% | 5 | 17.9% | 8 | 10.3% |
| Constipation | 0 | 47 | 94.0% | 24 | 85.7% | 71 | 91.0% |
| | 1 | 3 | 6.0% | 4 | 14.3% | 7 | 9.0% |
| Diarrhoea | 0 | 40 | 80.0% | 24 | 85.7% | 64 | 82.1% |
| | 1 | 6 | 12.0% | 0 | 0.0% | 6 | 7.7% |
| | 2 | 3 | 6.0% | 4 | 14.3% | 7 | 9.0% |
| | 3 | 1 | 2.0% | 0 | 0.0% | 1 | 1.3% |
| Faecal incontinence | 0 | 23 | 46.0% | 9 | 32.1% | 32 | 41.0% |
| | 1 | 13 | 26.0% | 6 | 21.4% | 19 | 24.4% |
| | 2 | 14 | 28.0% | 12 | 42.9% | 26 | 33.3% |
| | 3 | 0 | 0.0% | 1 | 3.6% | 1 | 1.3% |
| Flatulence | 0 | 20 | 40.0% | 11 | 39.3% | 31 | 39.7% |
| | 1 | 19 | 38.0% | 12 | 42.9% | 31 | 39.7% |
| | 2 | 11 | 22.0% | 5 | 17.9% | 16 | 20.5% |
| Frequency | 0 | 27 | 54.0% | 11 | 39.3% | 38 | 48.7% |
| | 1 | 11 | 22.0% | 11 | 39.3% | 22 | 28.2% |
| | 2 | 11 | 22.0% | 6 | 21.4% | 17 | 21.8% |
| | 3 | 1 | 2.0% | 0 | 0.0% | 1 | 1.3% |
| Rectal bleeding | 0 | 23 | 46.0% | 12 | 42.9% | 35 | 44.9% |
| | 1 | 17 | 34.0% | 11 | 39.3% | 28 | 35.9% |
| | 2 | 8 | 16.0% | 5 | 17.9% | 13 | 16.7% |
| | 3 | 2 | 4.0% | 0 | 0.0% | 2 | 2.6% |
| Rectal pain | 0 | 41 | 82.0% | 22 | 78.6% | 63 | 80.8% |
| | 1 | 8 | 16.0% | 5 | 17.9% | 13 | 16.7% |
| | 2 | 1 | 2.0% | 1 | 3.6% | 2 | 2.6% |
| Tenesmus | 0 | 21 | 42.0% | 11 | 39.3% | 32 | 41.0% |
| | 1 | 17 | 34.0% | 9 | 32.1% | 26 | 33.3% |
| | 2 | 12 | 24.0% | 8 | 28.6% | 20 | 25.6% |
| Urgency | 0 | 10 | 20.0% | 5 | 17.9% | 15 | 19.2% |
| | 1 | 20 | 40.0% | 9 | 32.1% | 29 | 37.2% |
| | 2 | 20 | 40.0% | 14 | 50.0% | 34 | 43.6% |
| GI disorders | 0 | 48 | 96.0% | 27 | 96.4% | 75 | 96.2% |
| | 1 | 2 | 4.0% | 1 | 3.6% | 3 | 3.8% |

A4.3 12 month CTCAE toxicities

| | | Control N=26 | | HBO N=46 | | Total N=72 | |
|---------------------|---|-----------------|-------|-------------|-------|---------------|-------|
| | | N | % | HBO | % | N | % |
| Abdominal pain | 0 | 14 | 53.8% | 20 | 43.5% | 34 | 47.2% |
| | 1 | 7 | 26.9% | 12 | 26.1% | 19 | 26.4% |
| | 2 | 1 | 3.8% | 9 | 19.6% | 10 | 13.9% |
| | 3 | 4 | 15.4% | 5 | 10.9% | 9 | 12.5% |
| Bloating | 0 | 12 | 46.2% | 22 | 47.8% | 34 | 47.2% |
| | 1 | 10 | 38.5% | 20 | 43.5% | 30 | 41.7% |
| | 2 | 4 | 15.4% | 4 | 8.7% | 8 | 11.1% |
| Constipation | 0 | 25 | 96.2% | 43 | 93.5% | 68 | 94.4% |
| | 1 | 1 | 3.8% | 2 | 4.3% | 3 | 4.2% |
| | 2 | 0 | 0.0% | 1 | 2.2% | 1 | 1.4% |
| Diarrhoea | 0 | 23 | 88.5% | 39 | 84.8% | 62 | 86.1% |
| | 1 | 1 | 3.8% | 1 | 2.2% | 2 | 2.8% |
| | 2 | 0 | 0.0% | 6 | 13.0% | 6 | 8.3% |
| | 3 | 2 | 7.7% | 0 | 0.0% | 2 | 2.8% |
| Faecal incontinence | 0 | 10 | 38.5% | 18 | 39.1% | 28 | 38.9% |
| | 1 | 5 | 19.2% | 14 | 30.4% | 19 | 26.4% |
| | 2 | 10 | 38.5% | 13 | 28.3% | 23 | 31.9% |
| | 3 | 1 | 3.8% | 1 | 2.2% | 2 | 2.8% |
| Flatulence | 0 | 10 | 38.5% | 16 | 34.8% | 26 | 36.1% |
| | 1 | 11 | 42.3% | 15 | 32.6% | 26 | 36.1% |
| | 2 | 5 | 19.2% | 15 | 32.6% | 20 | 27.8% |
| Frequency | 0 | 12 | 46.2% | 17 | 37.0% | 29 | 40.3% |
| | 1 | 7 | 26.9% | 13 | 28.3% | 20 | 27.8% |
| | 2 | 6 | 23.1% | 16 | 34.8% | 22 | 30.6% |
| | 3 | 1 | 3.8% | 0 | 0.0% | 1 | 1.4% |
| Rectal bleeding | 0 | 11 | 42.3% | 25 | 54.3% | 36 | 50.0% |
| | 1 | 13 | 50.0% | 14 | 30.4% | 27 | 37.5% |
| | 2 | 2 | 7.7% | 6 | 13.0% | 8 | 11.1% |
| | 3 | 0 | 0.0% | 1 | 2.2% | 1 | 1.4% |
| Rectal pain | 0 | 20 | 76.9% | 35 | 76.1% | 55 | 76.4% |
| | 1 | 3 | 11.5% | 11 | 23.9% | 14 | 19.4% |
| | 2 | 1 | 3.8% | 0 | 0.0% | 1 | 1.4% |
| | 3 | 2 | 7.7% | 0 | 0.0% | 2 | 2.8% |
| Tenesmus | 0 | 17 | 65.4% | 20 | 43.5% | 37 | 51.4% |
| | 1 | 4 | 15.4% | 15 | 32.6% | 19 | 26.4% |
| | 2 | 5 | 19.2% | 11 | 23.9% | 16 | 22.2% |
| Urgency | 0 | 6 | 23.1% | 13 | 28.3% | 19 | 26.4% |
| | 1 | 8 | 30.8% | 5 | 10.9% | 13 | 18.1% |
| | 2 | 11 | 42.3% | 24 | 52.2% | 35 | 48.6% |
| | 3 | 1 | 3.8% | 4 | 8.7% | 5 | 6.9% |
| GI disorders | 0 | 22 | 84.6% | 42 | 91.3% | 64 | 88.9% |
| | 1 | 1 | 3.8% | 2 | 4.3% | 3 | 4.2% |
| | 2 | 3 | 11.5% | 2 | 4.3% | 5 | 6.9% |